

HOMEWRECKERS

Stars disturbing their ancestral homes

Alyssa Goodman* (*Harvard-Smithsonian Center for Astrophysics*)

Héctor Arce (*Yale University Astronomy Department*)

Michelle Borkin* (*Harvard School of Engineering and Applied Science*)

Michael Halle* (*Brigham & Women's Hospital*)

Jens Kauffmann* (*JPL*)

Jaime Pineda (*Harvard-Smithsonian Center for Astrophysics*)


Sanjana Sharma (*Winsor School/Harvard-Smithsonian CfA*)

Lawrence Valverde (*Harvard College*)

*=*also Harvard Initiative in Innovative Computing*

Motivation

“Bipolar Flows”




cf. Matzner & McKee...; Nakamura & Li 2007

“Shells”

HOMEWRECKERS?

cf. Quillen et al. 2005; Churchwell et al...; Beaumont & Williams 2009


What Stars can do to the ISM



20 cm VLA from MAGPIS (Helfand et al. 2006) & MIR from Spitzer GLIMPSE (see Churchwell et al.)
3.6, 4.5, 8.0, 20cm (Luptonized, see Lupton et al. 2004)
image "height" is 1.6 degrees (e.g. 140 pc at 5 kpc)




Evolution of an HII Region in a Turbulent Medium



from S.J.Arthur 2007

M17




Hubble Image of the Swan Nebula
Photograph courtesy of NASA/STSCI/Jeff Hester, Arizona State University




© 2007 National Geographic Society. All rights reserved.

This is not real..but it could be.



Synthetic [OIII], Ha and [NII] emission-line image from a 512^3 numerical simulation: Mellema, Henney, Arthur & Vázquez-Semadeni 2009



Is it only O stars & HII regions that “matter”?



IMF $O:B:A:F$ populations would be: $1:50:300:750$.




Field $O:B:A:F$ populations: $1:4e3:2e4:1e5$.

r-Oph is a B*
 (and it's NOT in the "r-Oph" Cluster!)



see Schnee, Ridge, Goodman & Li 2005.

Ionized Gas in the Ophiuchus Smoke Shell




SHASSA Data courtesy of John Gaustad

COMPLETE =

COordinated MOlecular PRobe LIne EXtinction Thermal
Emission Survey of Star-Forming Regions

Perseus



COMPLETE Collaborators,
2010:

Alyssa A. Goodman (CfA/IIC)

João Alves (Vienna)

Héctor Arce (Yale)

Michelle Borkin (Harvard SEAS/IIC)

Paola Caselli (Leeds, UK)

James DiFrancesco (HIA, Canada)

Jonathan Foster (B.U.)

Mark Heyer (UMASS/FCRAO)

Doug Johnstone (HIA, Canada)

Jens Kauffmann (JPL/Caltech)

Helen Kirk (CfA)

Di Li (JPL/Caltech)

Stella Offner (CfA)

Jaime Pineda (CfA, PhD Student)

Thomas Robitaille (CfA)

Erik Rosolowsky (UBC Okanagan)

Rahul Shetty (ITA Heidelberg)

Scott Schnee (HIA Victoria)



Mario Tafalla (OAN, Spain)



What B* HD 278942 Does to Perseus

Total Dust **Column Density** (0 to 15 mag A_V)
(Based on 60/100 microns)

Dust **Column Temperature** (25 to 45 K)
(Based on 60/100 microns)




Cold Molecular Cloud, Warm Shell

Column
Density

Column
Temperature

+

=



Note: see Shetty et al. 2010 for “column temperature” discussion

Shell is “behind,” but touching, Perseus Molecular Cloud

IRAS N_{dust}

H α

2MASS/NICER
Extinction


*

H- α emission, WHAM/SHASSA Surveys (see Finkbeiner 2003)


“What about Magnetic Fields?”

Column
Density

Column
Temperature




+





=

Overlays from *Ridge et al. 2006*, showing polarizaiton (magnetic field?) deeply influenced by shell







And, the shell effects CO/dust ratio too...




Goodman, Pineda & Schnee 2009

COMPLETE Perseus


Image size: 1305 x 733
VL: 63 WW: 127

-  mm peak (Enoch et al. 2006)
-  sub-mm peak (Hatchell et al. 2005, Kirk et al. 2006)
-  ^{13}CO (Ridge et al. 2006)
-  mid-IR IRAC composite from c2d data (Foster, Laakso, Ridge, et al. in prep.)
-  Optical image (Barnard 1927)



m: 1/249
Zoom: 227% Angle: 0






Perseus

3D Viz made with VolView

AstronomicalMedicine@iig

COMPLETE



Bipolar Outflows in Perseus

THE ASTROPHYSICAL JOURNAL, 715:1170–1190, 2010 June 1

© 2010. The American Astronomical Society. All rights reserved. Printed in the U.S.A.

doi:10.1088/0004-637X/715/2/1170



THE COMPLETE SURVEY OF OUTFLOWS IN PERSEUS

HÉCTOR G. ARCE¹, MICHELLE A. BORKIN², ALYSSA A. GOODMAN³, JAIME E. PINEDA³, AND MICHAEL W. HALLE^{4,5}

¹ Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520, USA; hector.arce@yale.edu

² School of Engineering and Applied Sciences, Harvard University, 29 Oxford Street, Cambridge, MA 02138, USA

³ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁴ Surgical Planning Laboratory, Department of Radiology, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115, USA

⁵ Initiative in Innovative Computing, Harvard University, 60 Oxford Street, Cambridge, MA 02138, USA

Received 2009 October 7; accepted 2010 April 9; published 2010 May 7

ABSTRACT

We present a study on the impact of molecular outflows in the Perseus molecular cloud complex using the COMPLETE Survey large-scale $^{12}\text{CO}(1-0)$ and $^{13}\text{CO}(1-0)$ maps. We used three-dimensional isosurface models generated in right ascension–declination–velocity space to visualize the maps. This rendering of the molecular line data allowed for a rapid and efficient way to search for molecular outflows over a large ($\sim 16 \text{ deg}^2$) area. Our outflow-searching technique detected previously known molecular outflows as well as new candidate outflows. Most of these new outflow-related high-velocity features lie in regions that have been poorly studied before. These new outflow candidates more than double the amount of outflow mass, momentum, and kinetic energy in the Perseus cloud complex. Our results indicate that outflows have significant impact on the environment immediately surrounding localized regions of active star formation, but lack the energy needed to feed the observed turbulence in the *entire* Perseus complex. This implies that other energy sources, in addition to protostellar outflows, are responsible for turbulence on a global cloud scale in Perseus. We studied the impact of outflows in six regions with active star formation within Perseus of sizes in the range of 1–4 pc. We find that outflows have enough power to maintain the turbulence in these regions and enough momentum to disperse and unbind some mass from them. We found no correlation between outflow strength and star formation efficiency (SFE) for the six different regions we studied, contrary to results of recent numerical simulations. The low fraction of gas that potentially could be ejected due to outflows suggests that additional mechanisms other than cloud dispersal by outflows are needed to explain low SFEs in clusters.



Key words: ISM: clouds – ISM: individual objects (Perseus) – ISM: jets and outflows – ISM: kinematics and dynamics – stars: formation – turbulence

Online-only material: color figures




“CPOCs”

COMPLETE Perseus Outflow Candidates

Note: I did not make up that name!



“CPOCs”



Perseus

Bipolar

Outflows

Arce et al. 2010a

Table 5
Physical Parameters of Active Star-forming Regions in Perseus

Name	M_{reg}^a (M_{\odot})	R_{reg}^b (pc)	Δv^c (km s^{-1})	T_{ex}^d (K)	v_{esc}^e (km s^{-1})	E_{grav}^f (10^{46} erg)	E_{turb}^g (10^{45} erg)	t_{diss}^h (10^5 yr)	L_{turb}^i (10^{32} erg s^{-1})
L1448	150	0.6	1.9	10	1.5	0.3	2.9	2.6	3.6
NGC 1333	1100	2.0	2.2	13	2.2	5.2	28.8	5.7	15.9
B1-Ridge	210	0.7	1.9	13	1.6	0.5	4.1	3.1	4.1
B1	430	0.9	2.1	13	2.0	1.8	10.2	2.9	11.2
B5	420	1.4	1.5	12	1.6	1.1	5.1	7.6	2.1
IC 348	620	0.9	1.8	15	2.4	3.7	10.9	3.0	11.4

Notes.

- ^a Mass of star-forming region, obtained using the procedure described in Section 5.1.
- ^b Radius estimate of the region obtained from the geometric mean of minor and major axes of the extent of the ^{13}CO integrated intensity emission.
- ^c Average velocity width (FWHM) of the $^{13}\text{CO}(1-0)$ line in the region.
- ^d Average excitation temperature of region.
- ^e Escape velocity, given by $\sqrt{2GM_{\text{reg}}/R_{\text{reg}}}$.
- ^f Gravitational binding energy given by $GM_{\text{reg}}^2/R_{\text{reg}}$.
- ^g Turbulence energy given by $\frac{3}{16ln2}M_{\text{reg}}\Delta v^2$.
- ^h Turbulence dissipation time, see Section 5.2.1.
- ⁱ Turbulence energy dissipation rate given by $E_{\text{turb}}/\tau_{\text{diss}}$.

Table 6
Total Outflow Mass, Momentum, Energy, and Luminosity in Star-forming Regions

Name	M_{flow}^a (M_{\odot})	P_{flow}^a ($M_{\odot} \text{ km s}^{-1}$)	E_{flow}^a (10^{44} erg)	L_{flow}^b (10^{32} erg s^{-1})
L1448	1.0/5	3.1/21.7	1.2/12	8
NGC 1333	5.0/25	17.4/121.8	6.9/69	44
B1-Ridge	1.1/5.5	3.2/22.4	1.0/10	6
B1	1.5/7.5	6.2/43.4	3.1/31	20
IC 348	4.2/21	7.7/53.9	1.5/15	10
B5	12.8/64	22.3/156.1	4.1/41	26

Notes.

- ^a Values before and after the slash are the original estimates and the estimates adjusted by the correction factor, respectively (see Section 5.1).
- ^b Outflow luminosity, $L_{\text{flow}} = E_{\text{flow}}/\tau_{\text{flow}}$, obtained using the value of the total outflow kinetic energy adjusted by the correction factor and using an average outflow timescale of 5×10^4 yr.


Table 7
Quantitative Assessment of Outflow Impact on Star-forming Regions

Name	$E_{\text{flow}}/E_{\text{turb}}$	$r_L = L_{\text{flow}}/L_{\text{turb}}$	$E_{\text{flow}}/E_{\text{grav}}$	M_{esc}^a (M_{\odot})	$M_{\text{esc}}/M_{\text{reg}}$
L1448	0.41	2.1	0.40	15	0.10
NGC 1333	0.30	3.4	0.17	76	0.07
B1-Ridge	0.24	1.5	0.20	14	0.07
B1	0.30	1.7	0.17	21	0.05
IC 348	0.14	0.8	0.04	23	0.04
B5	0.80	12.4	0.37	98	0.23

Note. ^a Escape mass, given by $M_{\text{esc}} = P_{\text{out}}/v_{\text{esc}}$ (see Section 5.2.3).

Typically 20% binding energy in flows.

Bottom line
local influence significant,
HOMEWRECKERS **not.**



Properties of Molecular Clouds as “Equivalent Momentum” (using Larson 1981)

grey boxes mark lines of constant “momentum,” as labeled

Table 6

Total Outflow Mass, Momentum, Energy, and Luminosity in Star-forming Regions

Name	M_{flow}^a (M_{\odot})	P_{flow}^a ($M_{\odot} \text{ km s}^{-1}$)	E_{flow}^a (10^{44} erg)	L_{flow}^b ($10^{32} \text{ erg s}^{-1}$)
L1448	1.0/5	3.1/21.7	1.2/12	8
NGC 1333	5.0/25	17.4/121.8	6.9/69	44
B1-Ridge	1.1/5.5	3.2/22.4	1.0/10	6
B1	1.5/7.5	6.2/43.4	3.1/31	20
IC 348	4.2/21	7.7/53.9	1.5/15	10
B5	12.8/64	22.3/156.1	4.1/41	26

Notes.

^a Values before and after the slash are the original estimates and the estimates adjusted by the correction factor, respectively (see Section 5.1).

^b Outflow luminosity, $L_{\text{flow}} = E_{\text{flow}}/\tau_{\text{flow}}$, obtained using the value of the total outflow kinetic energy adjusted by the correction factor and using an average outflow timescale of $5 \times 10^4 \text{ yr}$.

Roughly true statement


Simulations show that ~kinetic energy observed must be injected every crossing time to maintain turbulence.

For reference: crossing time $\sim 10 \text{ pc}/2 \text{ km s}^{-1} = 5 \text{ Myr}$;

“flow time” = 0.05 Myr , so

flows per crossing time = $5 \text{ Myr}/0.05 \text{ Myr} = 100$


See also Curtis et al. (6/10 astro-ph) for a targeted study (*not a full census*)






“Shells”

Perseus Outflows & Shells



- | | | | | | |
|---|--------------------|---|---------------------------|---|-----------------------|
| + | Red Shifted CPOCs | ○ | New outflows | ◇ | IRAS Sources |
| + | Blue Shifted CPOCs | ○ | Known outflows | ◇ | Known Outflow Sources |
| × | HH Objects | ○ | Many small known outflows | ○ | “Shells” |
| | | ○ | Outflow extensions | | |

Arce et al. 2010 a,b




Perseus

3D Viz made with VolView

AstronomicalMedicine@iig




COMPLETE

Perseus Shells in ^{13}CO




Perseus Shells in Spitzer MIPS 24 mm

(Images from Spitzer c2d: Rebull et al. 2007)




“Cinema Arce”

x: 100	y: 523	z: 296	value: 0.312653	K
Ra 03h 48m 28.082s	Dec 33d 20m 34.95s	Vel: 8.83 km/s		




x: 168	y: 150	z: 257	value: -0.554963
Ra 03h 41m 19.851s	Dec 31d 55m 51.95s	Vel: 6.35 km/s	



IRAS 03382+3145

Shells in Perseus



10x momentum & energy of bipolar flows (now)

*mass has been multiplied by 2.5 to estimate opacity correction (full correction coming soon!)
also, IC348/Omicron Per HII region is not included, yet*

Perseus

Arce et al. 2010a,b

OBSERVED Momentum in **Shells** is $\sim 10\times$ **more** than in **bipolar flows**...

“Bipolar” Flows

Shells

Table 6

Total Outflow Mass, Momentum, Energy, and Luminosity in Star-forming Regions

Name	M_{flow}^a (M_{\odot})	P_{flow}^a ($M_{\odot} \text{ km s}^{-1}$)	E_{flow}^a (10^{44} erg)	L_{flow}^b (10^{32} erg s^{-1})
L1448	1.0/5	3.1/21.7	1.2/12	8
NGC 1333	5.0/25	17.4/121.8	6.9/69	44
B1-Ridge	1.1/5.5	3.2/22.4	1.0/10	6
B1	1.5/7.5	6.2/43.4	3.1/31	20
IC 348	4.2/21	7.7/53.9	1.5/15	10
B5	12.8/64	22.3/156.1	4.1/41	26

Notes.

^a Values before and after the slash are the original estimates and the estimates adjusted by the correction factor, respectively (see Section 5.1).

^b Outflow luminosity, $L_{\text{flow}} = E_{\text{flow}}/\tau_{\text{flow}}$, obtained using the value of the total outflow kinetic energy adjusted by the correction factor and using an average outflow timescale of 5×10^4 yr.


Name	Mass	Vexp	Momentum	Energy	Driver
B5-IRS4	62	2	124	2.5	IRAS 03446+3254 (B5-IRS4)
IC348 Shells	42	1	42	0.4	Dust Bowl, HD281159, and J03442106+32073862MASS
IRAS03390+3158	45	5	225	11.1	IRAS 03390+3158
IRAS03382+3145	181	2	362	7.2	IRAS 03382+3145
HD278942	89	5	446	22.2	HD278942
DR-B1 Shell	49	5	244	12.1	SSTcd2 J033525.4+310925
NGC1333-SW	40	2	80	1.6	multiple candidates
NGC1333-NW	126	2	252	5	multiple candidates
Totals	634		1775	62.1	


130 M_{\odot}
 420 $M_{\odot} \text{ km s}^{-1}$
 1.8×10^{46} erg

$\times 10 =$

1600 M_{\odot}
 4500 $M_{\odot} \text{ km s}^{-1}$
 16×10^{46} erg

For reference: the shell surrounding B-star **HD 278942** (G159.6-18.5; Ridge et al. 2006) is found to have a momentum of approximately **1400 $M_{\odot} \text{ km s}^{-1}$** and **$5 \times 10^{46}$ ergs of energy.**





Properties of Molecular Clouds
as
“Equivalent Momentum”
(using Larson 1981)

grey boxes mark lines of constant
“momentum,” as labeled

Are “upshifts” needed or correct?

Note theory gives ~ 10 to $1000 M_{\odot} \text{ km s}^{-1}$ per B-star wind.

Let's think about the basics...

(Underlined quantities are “measured”...others are “derived” or estimated.)

To calculate “**work**” done by flows, we need

$$\text{work} = (\text{force} * \text{distance}) \quad [\text{ergs} = \text{gm cm}^2 \text{ s}^{-2}]$$

$$\text{force of flow} = (\text{momentum}/\text{time}) \quad [\text{gm cm s}^{-2}]$$

$$\text{instantaneous momentum} = \text{mass} * \text{velocity} \quad [\text{gm cm s}^{-1}]$$

$$\text{mechanical luminosity} = \text{“kinetic energy”}/\text{time} \quad [\text{erg s}^{-1}]$$

Note that work has units of energy but is not necessarily $= (1/2) mv^2$.

For a nice analytic treatment, see Matzner 2007.

How much do (spherical) winds matter more generally?

Why should we even think about this?
(Perseus HD 278942 example Ridge et al. 2006; Arce, Borkin, Pineda, Goodman 2010a,b)

What **energy/momentum would be needed** to drive “turbulence”?
(Larson plot, analytic ideas, simulations)

How many (B) stars are available in “relevant” volumes?
(Sharma Catalog)

What **could/do (B) stars** do?
(Valverde Thesis)

Where, exactly are stars with winds, and **when** (for how long)?
(very hard to deduce from observations!)

What is **net effect** of stellar winds in **context** to e.g. bipolar outflows, HII regions?
(HII regions >> shells >= bipolar outflows)

What are the biggest **uncertainties** in this story?
(“net” momentum/energy deposition for various stellar types, **space-time-evolution** of winds)

Differential Diagnosis

- ▶ **Spitzer** Data suggests **shells** in **massive SFRs** (...but see *Beaumont & Williams 2009*); **COMPLETE** data suggests several **shells** in **wimpier** locales
- ▶ **Theorists** suggest spherical “mass-loaded winds” from massive stars don’t matter, b/c **radiation** is **dominant**.
- ▶ **Length & time** scales effected by shells may be **different** (larger & longer) than for bipolar outflows.
- ▶ Recent **numerical** work by Wang et al., Nakamura & Li, makes **outflows** look better (more **effective**) than Perseus data do (Arce et al. 2010). Why?
- ▶ **Time/spatial** scales + **coupling** are not well-known.
- ▶ We need a way to learn patient **histories**, as “**all patients lie.**”

^{13}CO is in the
“middle”

What is “Seamless Astronomy”?

^{12}CO is at the
“edges”



Extra Slides from Ringberg 2010

Sanjana Sharma's new Catalog (10,000 B-stars w/in 600 pc)
+
Lawrence Valverde's Harvard Senior Thesis
(theoretical/observational comparison for B-stars like
HD278942)

Preliminary Conclusion:


*Plausible that B-stars (and maybe less massive stars too!?) make
key contribution where they are most massive stars around.*

Goodman et al. 2011


For various conditions, we need to estimate:

○ Bipolar Flows ○ “Shells” ○ HII Regions

on pc
scales




on 10 pc
scales




etc...

IRAC and MIPS composite image from Spitzer Space Telescope website:
Color image made with: 4.5 μm (blue); 8.0 μm (green); 24 μm (red).

Color circles show position of shells shown in SnapZ movie. Note that the circles are included just to show the approximate positions of the shells (the size of the circle does not mean anything). The shells shown with the red and the blue circles were discovered using the Spitzer images (and are a bit hard to see in the 12CO channel maps). The yellow circle denotes a shell that was found in the 12CO channel maps, but it is not seen in the Spitzer images.




x: 244	y: 367	z: 280	value: 1.06695	K
Ra 03h 44m 8.183s	Dec 32d 20m 46.95s	Vel: 7.81 km/s		



Where do shells go on Larson's diagram?

*...but, see Beaumont & Williams 2009



HST view of NGC 604 in M33 (1 Mpc away)





Figure 3.9: Apparent B Star Shell in Glimpse Data
The yellow circle surrounds an example of coincidental overlap between B stars and molecular cloud structure separated radial to our line of sight by hundreds of parsecs.



A Venn diagram consisting of two overlapping circles. The left circle is olive green and contains the text 'B Star Shells'. The right circle is teal and contains the text 'Molecular Clouds'. The overlapping area in the center is a lighter shade of green and contains a white question mark. The background is black.

B
Star
Shells

Molecular
Clouds

?

World Wide Telescope

OB Associations in Orion B Stars in Orion's Belt





Table 3.2: Variables for Molecular Cloud – B Star Shell Intersection

Variable	Meaning	Estimated Value
R_s	Radius of one B star shell	3 pc
V_s	Volume of one B star shell	$1.1 \times 10^2 \text{ pc}^3$
N	Number of B stars within volume, V_{space}	10^4
V_{space}	Volume of catalogue region	$1.7 \times 10^8 \text{ pc}^3$
V_w	Volume of space for ff	$2.4 \times 10^{10} \text{ pc}^3$
ff	Molecular cloud filling factor	8×10^{-3}
f	Fraction of B star shells located within Molecular Clouds ¹	$5 \times 10^{-5} - 8 \times 10^{-1}$

3.1 General Formulation

The effect of B stars on turbulence in molecular clouds can be inferred from the energy and momentum deposited by B stars into molecular clouds. For a given environment, this energy is a function over time of the number of B stars and the energy deposited into molecular clouds per star. This can be represented symbolically as

$$\dot{E} = N \dot{E}_d \quad (3.1)$$

where \dot{E} is the total energy imparted to a molecular cloud per unit time, N is the number of B stars within a given volume of space, and \dot{E}_d is a function giving the energy deposited by each B star into a molecular cloud per unit time. The function \dot{E}_d can be further defined as

$$\dot{E}_d = f \dot{E}_s \varepsilon \quad (3.2)$$

where f is the integrated volume of shells from B stars per volume of molecular clouds in the galaxy, \dot{E}_s is the energy output of one B star per unit time, and ε is the energy transfer efficiency from the B star winds to the molecular cloud. Finally, this formula must be integrated over the total B star lifetime for which winds are active. Thus, the final equation for which values must be found in order to determine the impact of B stars on molecular clouds is

$$E = \int_{InteractionStart}^{InteractionStop} (Nf\varepsilon \dot{E}_s) dt \quad (3.3)$$

How much energy/time?
How many stars?

What is geometric overlap?
(deposition)


Energy transfer efficiency?
(hardest part?)

Duration of interaction?

Since only the kinetic part of the energy is relevant in relation to turbulence, a similar relation in terms of momentum is more illuminating and can be defined as

$$MV_{\infty} = \int_{InteractionStart}^{InteractionStop} (Nf\mu \dot{M} V_{\infty S}) dt \quad (3.4)$$

where MV_{∞} is the momentum imparted to the interstellar medium, $\dot{M} V_{\infty S}$ is the momentum output per unit time per star, and μ is the momentum transfer efficiency from the B star winds to the molecular clouds.



$$ff = 0.0008$$

$$ff \times \frac{N \times V_s}{V_{space}} \approx 5.3 \times 10^{-5}$$


Volume of B Star Shells

≈ 0.8

Volume of Molecular Clouds



AG Comments: Differentiating between (near-star) effects of mass-loaded wind, and ultimate effects of (far-from-star) net momentum (mass-loaded + radiation-driven expanding heated shell) is hard... needs simulations....



Summary of Momentum Ranges for HD 278942

Method	Lower Bound ($M_A \text{ km s}^{-1}$)	Upper Bound ($M_A \text{ km s}^{-1}$)
$\dot{M} V_\infty = \eta \left(\frac{L_*}{c} \right)$	11	1100
$\dot{M} V_\infty = \dot{M} * (V_\infty / V_{esc}) * V_{esc}$	10	300
Observation		500

Momentum from
Mass-Loaded Winds

~10 – 500 M_{\odot} km/s

B Star Shell and
Molecular Cloud
Overlap

~ 5×10^{-5} – 0.8



Impact of B Star Shells per
Molecular Cloud

Momentum from
Mass-Loaded Winds

~10 – 500 M_A km/s

B Star Shell and
Molecular Cloud
Overlap

~ 5×10^{-5} – 0.8



Impact of B Star Shells per
Molecular Cloud

~ 5×10^{-4} – 400 M_A km/s

How Influential?

Relative Number	“Typical” Mass	Spectral Type	Wind Energy	Kinetic Energy	Lifetime (Myr)
1	20	0			6
50	4	B			300
300	1.8	A			2000
750	1.2	F			~10,000
>1500	0.9	G-K-M			>10,000

$$L \propto M^{3.5} \quad \tau_{ms} \approx 10^{10} \text{ years} \cdot \left[\frac{M}{M_{\odot}} \right] \cdot \left[\frac{L_{\odot}}{L} \right] = 10^{10} \text{ years} \cdot \left[\frac{M_{\odot}}{M} \right]^{2.5}$$

Microsoft WorldWide Telescope Web Client

http://www.worldwidetelescope.org/webclient/default.aspx?wtml=http%3a%2f%2fwww.worldwidetelescope.org% Google

Arxivsorter Open in Papers Mendeley Import QR46 AGraphX ADS Best facetedADS Mundie AG Home Page WWTSL Harvard IIC: Projects Wikis Etc. Google Calendar

Explore Guided Tours Search View Settings

Collections > Open Collections > M17-L-RGB FRC_jpg > 1 of 1

M17-L-RGB FRC_jr

M17

150 pc at 2 kpc

Look At Sky Imagery USNOB: US Naval Observatory Info Image Crossfade 1 of 5






Sagittarius Pluto Omega Nebula Omega Nebula NGC6554 NGC6561 Black Hole NGC6567

RA : 18h21m02s Dec : -16:13:06

Done

Perseus (Valverde Slide!)

Results of Ridge et al. 2006

-  mm peak (Enoch et al. 2006)
-  sub-mm peak (Hatchell et al. 2005, Kirk et al. 2006)
-  ^{13}CO (Ridge et al. 2006)
-  mid-IR IRAC composite from c2d data (Foster, Laakso, Ridge, et al. in prep.)
-  Optical image (Barnard 1927)

