

Introduction to Magnetohydrodynamics

Nick Murphy

Harvard-Smithsonian Center for Astrophysics

July 21, 2017



To MHD and beyond!

- ▶ What is MHD?
- ▶ The equations of MHD and their physical meaning
- ▶ Waves in MHD
 - ▶ Alfvén waves
 - ▶ Slow magnetosonic waves
 - ▶ Fast magnetosonic waves
- ▶ Beyond MHD
 - ▶ Extensions to MHD
 - ▶ Plasma kinetic theory
- ▶ Magnetic reconnection
- ▶ Final thoughts

What is MHD?

- ▶ **Fluid dynamics** studies how fluids behave in response to forces
 - ▶ How do rivers flow?
 - ▶ How do we breathe?
- ▶ **Electromagnetism** studies the effects of electric and magnetic fields
 - ▶ What forces are exerted on free protons and electrons?
 - ▶ How does light work?
- ▶ **Magnetohydrodynamics** couples Maxwell's equations of electromagnetism with fluid dynamics to describe the large-scale behavior of conducting fluids such as plasmas
 - ▶ How does plasma behave in the solar atmosphere and wind?
 - ▶ How can we use magnetic fields to confine plasma?

MHD is important in solar physics, astrophysics, space plasma physics, and in laboratory plasma experiments



Left: The International Thermonuclear Experimental Reactor (a tokamak currently under construction in France)

Right: The solar wind interacting with Earth's magnetosphere

MHD at a glance (SI units)

Continuity Equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{V}) = 0$$

Momentum Equation

$$\rho \left(\frac{\partial}{\partial t} + \mathbf{V} \cdot \nabla \right) \mathbf{V} = \mathbf{J} \times \mathbf{B} - \nabla p$$

Ampere's law

$$\mu_0 \mathbf{J} = \nabla \times \mathbf{B}$$

Faraday's law

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}$$

Ideal Ohm's law

$$\mathbf{E} + \mathbf{V} \times \mathbf{B} = 0$$

Divergence constraint

$$\nabla \cdot \mathbf{B} = 0$$

Adiabatic Energy Equation

$$\frac{d}{dt} \left(\frac{p}{\rho^\gamma} \right) = 0$$

Definitions: \mathbf{B} , magnetic field; \mathbf{V} , plasma velocity; \mathbf{J} , current density; \mathbf{E} , electric field; ρ , mass density; p , plasma pressure; γ , ratio of specific heats (usually 5/3); t , time.

The MHD approximation


- ▶ Assume the plasma behaves like a fluid
 - ▶ Macroscopic behavior (long timescales, large distances)
 - ▶ Maxwellian particle distributions
- ▶ Ignore the most significant physics advances since 1860:
 - ▶ Relativity ($v^2 \ll c^2$)
 - ▶ Quantum mechanics
 - ▶ Displacement current in Ampere's law
- ▶ Assume the plasma is fully ionized
 - ▶ Limited applicability to weakly ionized plasmas like the photosphere and chromosphere
- ▶ Ignore resistivity, viscosity, thermal conduction, and radiative cooling in ideal MHD

Vector calculus refresher¹

- ▶ The **gradient** of f (denoted by ∇f) is a vector pointing in the direction of the steepest slope of f . The magnitude of the gradient vector is the steepness of the slope.
- ▶ The **divergence** of \mathbf{F} (denoted by $\nabla \cdot \mathbf{F}$) is the extent to which there is more of a quantity exiting a small region in space than entering it.
- ▶ The **curl** of \mathbf{F} (denoted by $\nabla \times \mathbf{F}$) represents the swirliness of a vector field.

¹Adapted partially from Wikipedia

The continuity equation describes conservation of mass



- ▶ The **continuity equation** written in conservative form is:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{V}) = 0$$

- ▶ The partial derivative $\partial \rho / \partial t$ refers to the change in density at a single point in space
- ▶ The divergence of the mass flux $\nabla \cdot (\rho \mathbf{V})$ says how much plasma goes in and out of the region
- ▶ Put sources and sinks of mass on RHS

The second golden rule of astrophysics



*“The density of wombats
times the velocity of wombats
gives the flux of wombats.”*

The momentum equation is analogous to $ma = \mathbf{F}$

- ▶ The **momentum equation** is

$$\rho \left(\frac{\partial}{\partial t} + \mathbf{V} \cdot \nabla \right) \mathbf{V} = \mathbf{J} \times \mathbf{B} - \nabla p$$

Additional forces like gravity go on the right hand side.²

- ▶ The **total derivative** represents how much a quantity is changing as you follow a parcel of plasma:

$$\frac{D}{Dt} \equiv \frac{\partial}{\partial t} + \mathbf{V} \cdot \nabla$$


- ▶ Forces must cancel each other out in a static equilibrium:

$$\mathbf{J} \times \mathbf{B} = \nabla p$$

When $\mathbf{J} \times \mathbf{B} = 0$, the plasma is **force-free**

²If you neglect gravity, it may be your downfall! (I had to drop at least one pun in.)

The pressure gradient force $-\nabla p$ pushes plasma from regions of high pressure to low plasma pressure



The Lorentz force term includes two components

- ▶ The current density is given by the relative drift between ions and electrons:

$$\mathbf{J} = ne(\mathbf{V}_i - \mathbf{V}_e)$$

→ $\mathbf{J} \times \mathbf{B}$ is analogous to $\mathbf{F} = q\mathbf{V} \times \mathbf{B}$.


- ▶ Using vector identities and Ampere's law ($\mu_0 \mathbf{J} = \nabla \times \mathbf{B}$), we rewrite the Lorentz force term $\mathbf{J} \times \mathbf{B}$ as:

$$\mathbf{J} \times \mathbf{B} = \frac{(\mathbf{B} \cdot \nabla) \mathbf{B}}{\mu_0} - \nabla \left(\frac{B^2}{2\mu_0} \right)$$

However: the Lorentz force is orthogonal to \mathbf{B} , but these two terms are not.

The Lorentz force can be decomposed into two terms with forces orthogonal to \mathbf{B} using field line curvature

- ▶ The curvature vector κ points toward the center of curvature and gives the rate at which the tangent vector turns:




- ▶ We can then write the Lorentz force as

$$\underbrace{\mathbf{J} \times \mathbf{B}}_{\text{Lorentz force}} = \underbrace{\kappa \frac{B^2}{\mu_0}}_{\text{magnetic tension}} - \underbrace{\nabla_{\perp} \left(\frac{B^2}{2\mu_0} \right)}_{\text{magnetic pressure}} \quad (1)$$


where all terms are orthogonal to \mathbf{B} . The operator ∇_{\perp} takes the gradient only in the direction orthogonal to \mathbf{B} .

The magnetic tension force wants to straighten magnetic field lines



- ▶ The tension force is directed radially inward with respect to magnetic field line curvature

Regions of high magnetic pressure exert a force towards regions of low magnetic pressure



- ▶ The magnetic pressure is given by $p_B \equiv \frac{B^2}{2\mu_0}$

The ratio of the plasma pressure to the magnetic pressure is an important dimensionless number

- ▶ Define plasma β as

$$\beta \equiv \frac{\text{plasma pressure}}{\text{magnetic pressure}} \equiv \frac{p}{B^2/2\mu_0}$$

- ▶ If $\beta \ll 1$ then the magnetic field dominates
 - ▶ Solar corona
- ▶ If $\beta \gg 1$ then plasma pressure forces dominate
 - ▶ Solar interior
- ▶ If $\beta \sim 1$ then pressure/magnetic forces are both important
 - ▶ Solar chromosphere
 - ▶ Parts of the solar wind and interstellar medium
 - ▶ Some laboratory plasma experiments

Faraday's law tells us how the magnetic field varies with time

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}$$

But how do we get the electric field?

Ohm's law provides the electric field

- ▶ The ideal MHD Ohm's law is given by

$$\mathbf{E} + \mathbf{V} \times \mathbf{B} = 0$$

- ▶ In ideal MHD, the magnetic field is *frozen-in* to the plasma. If two parcels of plasma are connected by a magnetic field line at one time, then they will be connected by a magnetic field line at all other times.
- ▶ For resistive MHD, Ohm's law becomes

$$\mathbf{E} + \mathbf{V} \times \mathbf{B} = \eta \mathbf{J}$$

where η is the resistivity. Resistivity allows the frozen-in condition to be broken.

- ▶ Can also include the Hall effect which is important on short length scales


With Ohm's law we can rewrite Faraday's law as the induction equation

- ▶ Using the resistive Ohm's law:

$$\frac{\partial \mathbf{B}}{\partial t} = \underbrace{\nabla \times (\mathbf{V} \times \mathbf{B})}_{\text{convection}} + \underbrace{\frac{\eta}{\mu_0} \nabla^2 \mathbf{B}}_{\text{diffusion}}$$

Diffusion is usually represented by a second order spatial derivative.

- ▶ An example of resistive diffusion:



Thermal conduction is a common extension to MHD

- ▶ Heat diffuses much more quickly along magnetic field lines than perpendicular to them
 - ▶ Makes it more difficult to simulate plasmas
- ▶ The temperature along magnetic field lines is usually approximately constant
 - ▶ Exceptions: rapid heating events, rapid magnetic connectivity changes

Waves


- ▶ There are three primary waves that arise from MHD:
 - ▶ Alfvén wave
 - ▶ Slow magnetosonic wave
 - ▶ Fast magnetosonic wave
- ▶ There are two important speeds
 - ▶ The sound speed is given by

$$V_S \equiv \sqrt{\frac{\gamma p}{\rho}}$$

- ▶ The Alfvén speed is given by


$$V_A \equiv \frac{B}{\sqrt{\mu_0 \rho}}$$

Alfvén Waves



- ▶ Alfvén waves propagate at the Alfvén speed: $V_A \equiv \frac{B}{\sqrt{\mu_0 \rho}}$
- ▶ The restoring force is magnetic tension
- ▶ This is a shear wave with no compression involved
- ▶ Disturbances propagate parallel to **B**

Slow and Fast Magnetosonic Waves



- ▶ *Left:* The restoring forces for magnetosonic waves propagating perpendicular to \mathbf{B} are given by gas and magnetic pressure gradients. This shows a compressional wave.
- ▶ *Right:* The phase velocity of MHD waves are a function of angle when \mathbf{B} is in the z direction and β is small.
- ▶ Sound waves are magnetosonic waves propagating along \mathbf{B}




How useful is MHD?

- ▶ MHD is appropriate for large-scale, low-frequency behavior
- ▶ MHD is a good predictor of stability
 - ▶ But Non-MHD effects sometimes stabilize or destabilize...
- ▶ MHD is often inappropriate when there are non-Maxwellian distribution functions
 - ▶ Collisionless plasmas
 - ▶ Situations with lots of energetic, non-thermal particles
- ▶ MHD is a reasonable approximation for most solar physics applications, but effects beyond MHD are often important
- ▶ MHD is a mediocre description of laboratory plasmas



There are two general approaches to going beyond MHD

- ▶ **Extended MHD**
 - ▶ Keep the fluid approximation
 - ▶ Add more terms to the equations to include more effects
- ▶ **Kinetic theory**
 - ▶ Abandon the fluid approximation
 - ▶ Keep track of particle distribution functions
- ▶ Or . . . take both approaches simultaneously!

Magnetic Reconnection is the breaking and rejoining of magnetic field lines in a highly conducting plasma



Solar flares and CMEs are powered by magnetic reconnection



- ▶ Explosive release of magnetic energy
- ▶ Bidirectional Alfvénic jets
- ▶ Very efficient particle acceleration
- ▶ Flux ropes escape as coronal mass ejections (CMEs)

Magnetic reconnection is a fundamental process in laboratory and astrophysical plasmas

- ▶ Classical theories based on resistive diffusion predict slow reconnection (weeks to months...)
- ▶ Fast reconnection allows magnetic energy to be explosively converted into kinetic and thermal energy
- ▶ Collisionless or non-fluid effects are (probably) needed to explain why fast reconnection occurs in flares (tens of seconds to minutes!)

Summary

- ▶ MHD describes the macroscopic behavior of plasmas
- ▶ Each term in the MHD equations represents a different physical effect
- ▶ There are three types of MHD waves: Alfvén waves, fast magnetosonic waves, and slow magnetosonic waves
- ▶ Physics beyond MHD is often needed to describe plasma behavior
- ▶ Magnetic reconnection is the breaking and rejoining of magnetic field lines in a highly conducting plasma
 - ▶ Releases magnetic energy during solar flares and CMEs
 - ▶ Degrades confinement in laboratory plasmas

Useful references

- ▶ *The Physics of Plasmas* by T.J.M. Boyd and J.J. Sanderson. One of the most understandable introductions to plasma physics that I've found.
- ▶ *Magnetohydrodynamics of the Sun* by Eric Priest. Very useful resource for the mathematical properties of MHD as applied to the Sun.
- ▶ *Principles of Magnetohydrodynamics* by Hans Goedbloed and Stefaan Poedts. Good introduction to MHD with a broad focus on applications.
- ▶ *Ideal Magnetohydrodynamics* by Jeffrey Freidberg. Very good out-of-print introduction to MHD in particular. Later chapters focus more on laboratory plasmas.
- ▶ *Introduction to Plasma Physics and Controlled Fusion* by Francis Chen. A beginning graduate level introduction to plasma physics. Less emphasis on MHD.